arXiv:1703.04515v3 [math.SG] 21 Apr 2017

FUKAYA CATEGORIES OF PLUMBINGS AND MULTIPLICATIVE PREPROJECTIVE ALGEBRAS

TOLGA ETGÜ AND YANKI LEKILI

Abstract. Given an arbitrary graph Γ and non-negative integers g_v for each vertex v of Γ, let X_Γ be the Weinstein 4-manifold obtained by plumbing copies of $T^*\Sigma_v$ according to this graph, where Σ_v is a surface of genus g_v. We compute the wrapped Fukaya category of X_Γ (with bulk parameters) using Legendrian surgery extending our previous work [13] where it was assumed that $g_v = 0$ for all v and Γ was a tree. The resulting algebra is recognized as the (derived) multiplicative preprojective algebra (and its higher genus version) defined by Crawley-Boevey and Shaw [8]. Along the way, we find a smaller model for the internal DG-algebra of Ekholm-Ng [11] associated to 1-handles in the Legendrian surgery presentation of Weinstein 4-manifolds which might be of independent interest.

1. Introduction

In this paper, we continue our study from [13] of computing explicit models for the wrapped Fukaya category $\mathcal{W}(X_\Gamma)$ of the 4-dimensional open symplectic manifold X_Γ, obtained by plumbing cotangent bundles of surfaces according to a graph Γ. Previously in [13], we restricted to case of plumbings of cotangent bundles of 2-spheres and X_Γ was exhibited as a Legendrian surgery along a link $L_\Gamma \subset (S^3, \xi_{std})$ and the Chekanov-Eliashberg DG-algebra $\mathscr{B}_\Gamma$ of L_Γ was computed explicitly for any tree Γ. It is known that $\mathscr{B}_\Gamma$ is A_∞-quasi-isomorphic to the endomorphism algebra of the full subcategory of $\mathcal{W}(X_\Gamma)$ whose objects are the cocores of the 2-handles in the surgery presentation [5] (cf. [10, Thm. 2]). Furthermore, a combination of results of [5] and [1] implies that these cocores split-generate $\mathcal{W}(X_\Gamma)$. A proof of the stronger statement of generation (rather than split-generation) is ongoing work of two independent groups: Ganatra, Pardon, Shende, and Chantraine, Dimitroglou-Rizell, Ghiggini, Golovko. Thus, we take $\mathscr{B}_\Gamma$ as our primary object of interest.

We improve on the results of [13] in two aspects. Firstly, no restriction is imposed on Γ which was assumed to be a tree in [13] and only plumbings of T^*S^2's were considered. When Γ is not a tree or cotangent bundles of positive genus surfaces are used, Legendrian surgery description of X_Γ necessarily includes 1-handles, and the computation of $\mathscr{B}_\Gamma$ is more involved. More precisely, for any connected graph Γ with s vertices and $s + k - 1$ edges and $g = \sum_v g_v$, we shall construct a Legendrian L_Γ in $\#^{k+2g}(S^1 \times S^2, \xi_{std})$ with s components such that X_Γ is obtained by Legendrian surgery on L_Γ. A roadmap for an explicit computation of $\mathscr{B}_\Gamma$ is provided by the work of Ekholm-Ng [11] which involves a countably generated internal DG-algebra $\mathscr{I}$ associated to each 1-handle in the surgery presentation. We find a simplification by proving that a finitely generated DG-subalgebra of $\mathscr{I}$ is quasi-isomorphic to $\mathscr{I}$.

Secondly, when Γ is one of the A_n- and D_n-type trees (and conjecturally we also include E_6, E_7, E_8-types), it was shown in [13] that $\mathscr{B}_\Gamma$ is quasi-isomorphic to Ginzburg's DG-algebra $\mathscr{G}_\Gamma$, away from a small set of exceptional characteristics for the ground field $\mathbb{K}$. On the other hand, if Γ is a non-Dynkin tree, the explicit computation in [13] shows that $\mathscr{B}_\Gamma$ is a deformation of $\mathscr{G}_\Gamma$. In this case, it is also known that $\mathscr{G}_\Gamma$ is quasi-isomorphic to $H^0(\mathscr{G}_\Gamma) = \Pi_\Gamma$, the (additive) preprojective algebra attached to Γ (in contrast, $\mathscr{G}_\Gamma$ has non-trivial cohomology in every non-positive degree, when Γ is a Dynkin tree). It was argued that the associated formal deformation is trivial over a field of characteristic 0 since Π_Γ has no formal deformations as a Calabi-Yau algebra, but the precise determination of quasi-isomorphism type of $\mathscr{B}_\Gamma$ was not given.

In this paper, we first restrict to the case $g_v = 0$ and show that for any finite graph Γ, $\mathscr{B}_\Gamma$ is quasi-isomorphic to the derived multiplicative preprojective algebra $\mathscr{L}_\Gamma$ whose zeroth cohomology is isomorphic to the multiplicative preprojective algebra Λ_Γ defined in [8] by Crawey-Boevey and Shaw with motivation coming from the Deligne-Simpson problem. Next, we extend this to the case where g_v is arbitrary. The resulting DG-algebra is quasi-isomorphic to a positive genus version of the derived multiplicative algebra that appears in [3]. We shall denote this by $\mathscr{L}_{\Gamma,\mathbf{g}}$ and its zeroth cohomology $H^0(\mathscr{L}_{\Gamma,\mathbf{g}})$ by $\Lambda_{\Gamma,\mathbf{g}}$, where $\mathbf{g} = (g_1, g_2, \dots, g_s)$ is an s-tuple of non-negative integers recording the genus g_v associated to vertices of Γ. When $\mathbf{g}$ is omitted, it is understood that $g_v = 0$ for all v.

Before giving the definition of $\Lambda_{\Gamma,\mathbf{g}}$ and $\mathscr{L}_{\Gamma,\mathbf{g}}$, let us first introduce some notation. As in [13], we fix a field $\mathbb{K}$ and the semisimple ring

$$\mathbf{k} := \bigoplus_{v=1}^{s} \mathbb{K} e_v$$

where $e_v^2 = e_v$ and $e_v e_w = 0$ for $v \neq w$, and $\{1, \dots, s\}$ is the set of vertices of Γ. Here, we shall work over a bigger ring:

$$\mathbf{k}\langle \mathbf{t}^{\pm} \rangle = \mathbf{k}\langle t_1, t_1^{-1}, t_2, t_2^{-1}, \dots, t_s, t_s^{-1} \rangle / (t_v t_v^{-1} = t_v^{-1} t_v = e_v, t_v^{\pm} t_w^{\pm} = 0 \ \text{ for } \ v \neq w)$$

We also fix a spanning tree $T \subset \Gamma$ and orient the edges of Γ. The multiplicative preprojective algebra Λ_Γ is the associative $\mathbf{k}$-algebra obtained from $\mathbf{k}\langle \mathbf{t}^{\pm} \rangle$ by adjoining the generators

$$\begin{aligned} c_a, c_a^* &\ \text{ for any arrow } \ a \in \Gamma, \ \text{ and} \\ z_a, z_a^{-1} &\ \text{ for any arrow } \ a \in \Gamma \backslash T \\ \alpha_{v,i}, \alpha_{v,i}^{-1}, \beta_{v,i}, \beta_{v,i}^{-1} &\ \text{ for any vertex } \ v \in \Gamma \text{ and } i = 1, 2, \dots, g_v. \end{aligned}$$

(Note that there are indeed 4 generators associated to $a \in \Gamma \backslash T$: $c_a, c_a^*, z_a, z_a^{-1}$.)

The $\mathbf{k}\langle \mathbf{t}^{\pm} \rangle$-bimodule structure on Λ_Γ is obtained by extending the scalars from the $\mathbf{k}$-bimodule structure by

$$\begin{gathered} e_w c_a e_v = c_a, \quad e_v c_a^* e_w = c_a^*, \\ e_v z_a^{\pm} e_v = z_a^{\pm} \end{gathered}$$

if and only if $v = s(a)$ and $w = t(a)$ are the source and target vertices of the arrow a, respectively. The $\mathbf{k}$-bimodule structure on the remaining generators is the obvious one:

$$e_v \alpha_{v,i}^{\pm} e_v = \alpha_{v,i}^{\pm}$$
$$e_v \beta_{v,i}^{\pm} e_v = \beta_{v,i}^{\pm}$$

In all other cases, the product with an idempotent gives zero.

Note that the variables t_v are taken to be non-central (unless we specialise). The generators of Λ_Γ are subject to the following relations (all multiplications are read from right-to-left):

$$z_a z_a^{-1} = z_a^{-1} z_a = e_v, \quad \text{for } a \in \Gamma \backslash T \text{ and } v = s(a)$$
$$e_v + c_a^* c_a = z_a, \quad \text{for } a \in \Gamma \backslash T \text{ and } v = s(a)$$
$$\prod_{t(a)=v} (e_v + c_a c_a^*) = t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \prod_{s(a)=v} (e_v + c_a^* c_a), \quad \text{for } v \in \{1, \dots, s\}$$

where the terms in the products are ordered according a fixed order of all the arrows in Γ, and the empty product is understood to be the relevant idempotent. The notation $[x, y] = xyx^{-1}y^{-1}$ stands for the commutator.

It is known that $\Lambda_{\Gamma,\mathbf{g}}$ is independent of the orientation and order of the edges of Γ, and the choice of the spanning tree T (cf. [8, Thm. 1.4]). This also follows from our main result by the invariance properties of Chekanov-Eliashberg DG-algebra $\mathscr{B}_{\Gamma,\mathbf{g}}$.

A DG-algebra $\mathscr{L}_{\Gamma,\mathbf{g}}$ such that $H^0(\mathscr{L}_{\Gamma,\mathbf{g}}) \simeq \Lambda_{\Gamma,\mathbf{g}}$ is discussed in [3, Rem.5C & Ex.5.6] (see also [25]) under the name (universal) derived multiplicative preprojective algebra. In the current set-up, $\mathscr{L}_{\Gamma,\mathbf{g}}$ can be defined by adding generators with grading -1, in lieu of the last two relations above, to the set of generators of $\Lambda_{\Gamma,\mathbf{g}}$ (which are assigned grading 0) so that $H^0(\mathscr{L}_{\Gamma,\mathbf{g}}) = \Lambda_{\Gamma,\mathbf{g}}$. Namely, the differentials of these additional generators are

$$\partial \zeta_a = e_v + c_a^* c_a - z_a, \quad \text{for } a \in \Gamma \backslash T \text{ and } v = s(a)$$
$$\partial \tau_v = \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \prod_{s(a)=v} (e_v + c_a^* c_a), \quad \text{for } v \in \{1, \dots, s\}$$

We can now state our main theorem succinctly as follows:

Theorem 1. *For any graph Γ and $\mathbf{g}$, there is a quasi-isomorphism between the derived multiplicative preprojective algebra $\mathscr{L}_{\Gamma,\mathbf{g}}$ and the Chekanov-Eliashberg DG-algebra $\mathscr{B}_{\Gamma,\mathbf{g}}$.*

Remark 2. The main theorem is proved by identifying $\mathscr{B}_{\Gamma,\mathbf{g}}$ and $\mathscr{L}_{\Gamma,\mathbf{g}}$ by direct computation of the Chekanov-Eliashberg DG-algebra of the Legendrian $L_{\Gamma,\mathbf{g}}$. In view of the surgery results from [5], [10], our result provides computations of various wrapped Fukaya categories as:

$$\mathscr{B}_{\Gamma,\mathbf{g}} \cong \bigoplus_{v,w} CW^*(C_v, C_w)$$

where $CW^*(C_v, C_w)$ stands for the morphisms between the cocores C_v, C_w in the partially wrapped Fukaya of $(\#^{k+2g}(S^1 \times \mathbb{D}^3), \omega_{std})$ where all Legendrians L_v are used as stops

(see [10, Conj. 3]). In this setting, it is important that the indeterminates t_v's do not commute with the other generators. On the other hand, if one specialises to $t_v = e_v$ for $v = 1, \dots, s$, one recovers the fully wrapped Fukaya category of X_Γ where X_Γ is equipped with the exact symplectic form associated with its Weinstein structure. Setting the value of $\mathrm{Log}(t_v) = \omega([S_v])$ to other constants, has the effect of turning on non-exact (bulk) deformations of the symplectic form, by giving area to the core surfaces S_v of X_Γ. One can interpret this as a computation of bulk deformed wrapped Fukaya categories.

Remark 3. As in [10], let

$$\mathscr{A}_{\Gamma,\mathbf{g}} = \bigoplus_{v,w} CF^*(S_v, S_w)$$

be the endomorphism algebra of the compact exact Lagrangians in X_Γ which form the Lagrangian skeleton of the plumbing X_Γ. After letting $t_v = e_v$, for $v = 1, \dots, s$, one has a canonical augmentation

$$\epsilon : \mathscr{B}_{\Gamma,\mathbf{g}} \to \mathbf{k}$$

corresponding to the module $\bigoplus CF^*(C_v, S_v)$ in the (fully) wrapped Fukaya category of $\mathcal{W}(X_\Gamma)$, given explicitly by

$$\epsilon(c_a) = \epsilon(c_a^*) = 0 \ , \ \epsilon(\alpha_{v,i}^{\pm}) = \epsilon(\beta_{v,i}^{\pm}) = e_v = \epsilon(z_a^{\pm}), \ v = s(a)$$

By [10, Thm. 4], one can compute

$$\mathscr{A}_{\Gamma,\mathbf{g}} \simeq \mathrm{Rhom}_{\mathscr{B}_{\Gamma,\mathbf{g}}}(\mathbf{k}, \mathbf{k})$$

This could be useful in studying Fukaya categories of symplectic 4-manifolds. Namely, if one has any configuration of transversely intersecting compact Lagrangians in a symplectic 4-manifold whose intersection pattern is encoded by the decorated graph (Γ,$\mathbf{g}$), the corresponding A_∞-algebra is a deformation of $\mathscr{A}_{\Gamma,\mathbf{g}}$.

We also note that the grading in $\mathscr{B}_{\Gamma,\mathbf{g}}$ is supported in non-positive degrees. In higher dimensional plumbings, Abouzaid and Smith used this property and an additional requirement on the zeroth cohomology of cocores [2, Prop. 4.2], that guarantees that the only modules supported in a single degree are given by the compact cores with local systems, to deduce that the compact cores of the plumbing generate the compact Fukaya category. In our case, this additional requirement does not hold. Nonetheless, the core Lagrangians surfaces $\{S_v\}$ are known to generate the compact Fukaya category $\mathcal{F}(X_\Gamma)$ in several situations, including when Γ is of Dynkin type with $\mathbf{g} = 0$ ([21, Lem. 4.15] and [22, Cor. 5.8]). We also remark that Keating [16] gave examples of Weinstein 4-manifolds which are *partial compactifications* of plumbings, where an exact Lagrangian torus (equipped with certain local systems) is not generated by the core spheres.

Remark 4. After we wrote our first paper [13] on computing wrapped Fukaya categories of X_Γ, which studied the case where Γ is a tree and $\mathbf{g} = 0$, an interesting paper by Bezrukavnikov and Kapranov [3] appeared, where finite dimensional $\Lambda_{\Gamma,\mathbf{g}}$-modules are related to microlocal sheaves on the Lagrangian skeleton of X_Γ given by the union of transversely intersecting surfaces. In view of the correspondence between Fukaya categories of cotangent bundles and microlocal sheaves on the zero-section [17], Thm. (1) provides a

confirmation of the more general expectation of an equivalence between Fukaya categories of Weinstein manifolds and microlocal sheaves on (singular) Lagrangian skeleta for the manifolds X_Γ. This more general equivalence is the subject of ongoing work by Ganatra, Pardon and Shende.

Let us consider the case Γ is a tree and $\mathbf{g} = 0$, and specialise to $t_v = e_v$. By definition of $\mathscr{B}_\Gamma$, the differential $\partial\tau_v$ consists of words of (even) length ≥ 2. If we truncate it so that only words of length 2 appears, the resulting DG-algebra is known as the Ginzburg DG-algebra $\mathscr{G}_\Gamma$ which was studied in [13]. It is known that if Γ is not of Dynkin type, $\mathscr{G}_\Gamma$ has cohomology only in degree 0 (cf. Thm 8, Cor. 27 [13]) and $H^0(\mathscr{G}_\Gamma) = \Pi_\Gamma$ is the additive preprojective algebra. Now, by considering the word-length filtration, we get a spectral sequence from $\mathscr{G}_\Gamma$ to $\mathscr{B}_\Gamma$. At least when $\mathrm{char}\mathbb{K} = 0$, one can show that the length spectral sequence used in the above proof is degenerate. The reason for this was given in [13, pg. 7]. Namely, it is known that Π_Γ has no formal deformations as a Calabi-Yau algebra. However, the length spectral sequence does not converge strongly if Γ is non-Dynkin, hence one cannot conclude that $\mathscr{B}_\Gamma$ and $\mathscr{G}_\Gamma$ are quasi-isomorphic without having to complete both algebras. Indeed, for non-Dynkin Γ, the additive preprojective algebra $\Pi_\Gamma = H^0(\mathscr{G}_\Gamma)$ and the multiplicative preprojective algebra $\Lambda_\Gamma = H^0(\mathscr{B}_\Gamma)$ are not isomorphic. This can be seen by considering the representation varieties of Π_Γ and Λ_Γ - additive and multiplicative Nakajima quiver varieties. Thus, we deduce that our Conjecture 3 from [13] cannot hold. It is the (derived) multiplicative preprojective algebra $\mathscr{B}_\Gamma$ that is central to the study of wrapped Fukaya categories of plumbings, not the additive preprojective algebra (or its derived version).

Example 5. When Γ is just a single vertex labelled with g, then X_Γ is just the cotangent bundle of a genus g surface Σ_g. In this case, we see that the multiplicative preprojective algebra is simply the group-ring $\mathbb{K}[\pi_1(\Sigma_g)]$, which agrees with the based loop space homology of the cotangent fibre. This agrees with the computation given in [11].

Example 6. When Γ has a single vertex (labelled with 0) and a single loop, the symplectic manifold X_Γ obtained by self-plumbing of T^*S^2 was studied in detail in [20]. In this case, it is easy to see that (cf. [8, Ex. 1.3]) the multiplicative preprojective algebra (with $t = 1$) Λ_Γ is quasi-isomorphic to the commutative algebra $\mathbb{K}[x, y, (1 + xy)^{-1}]$, which agrees with the homological mirror symmetry statement proven in [20] (cf. [6]).

We end the introduction by pointing out an intriguing connection to (multiplicative) Nakajima quiver varieties that we plan to return to in the future. Note that such a connection was also observed in [3] (see also [23]) via calculations using microlocal sheaves.

It is well known that the moduli spaces of representations of the multiplicative preprojective algebra Λ_Γ has the structure of algebraic symplectic varieties known as the multiplicative Nakajima quiver varieties (studied by Yamakawa [24], see also Boalch [4]). By our result, any Lagrangian $L \subset X_\Gamma$ gives a representation $\bigoplus_v \mathrm{Hom}(C_v, L)$ of the multiplicative preprojective algebra $\bigoplus_{v,w} \mathrm{Hom}(C_v, C_w)$. Thus, our result can be interpreted as the statement that "moduli space of objects" in the derived Fukaya category of X_Γ can be identified with the corresponding multiplicative Nakajima quiver varieties. These are finite type varieties once one fixes a dimension vector consisting of $d_v \in \mathbb{Z}_{\geq 0}$ for each vertex

v of Γ and a stability condition encoded by $\theta_v \in \mathbb{Q}$ for each v, satisfying $\sum_v d_v\theta_v = 0$, and the values of $t_v \in \mathbb{C}^\times$. The choice of dimension vectors corresponds to fixing the rank $d_v = \dim_{\mathbb{K}}\mathrm{Hom}(L, C_v)$, and the choice of stability condition should correspond to fixing a choice of Bridgeland stability on the derived Fukaya category of X_Γ. Finally, note that the symplectic structures on the Nakajima quiver varieties correspond to the fact that $\mathscr{B}_{\Gamma,\mathbf{g}}$ is 2-Calabi-Yau (cf. [19]).

In Section 2, after reviewing the construction of Chekanov-Eliashberg algebra associated to a Legendrian surgery presentation of a Weinstein 4-manifold following Ekholm-Ng [11], we provide a simplification that results in a finitely generated DG-algebra even in the presence of 1-handles. In Section 3, we draw a Legendrian surgery presentation of plumbings of T^*S^2's according to an arbitrary graph and prove our main theorem for $\mathbf{g} = 0$. In Section 4, we extend the results of Section 3 to arbitrary $\mathbf{g}$.

Acknowledgments: T.E. is partially supported by the Technological and Research Council of Turkey through a BIDEB-2219 fellowship. Y.L. is supported in part by the Royal Society (URF) and the NSF grant DMS-1509141. We would like to thank Pavel Etingof, Lenhard Ng, and Ivan Smith for helpful correspondence.

2. Legendrian DG-algebra on a subcritical Weinstein 4-manifold

Recall that any Weinstein 4-manifold X has a Weinstein handlebody decomposition consisting of 0-, 1- and 2-handles. Assuming X is connected, one can find a handlebody decomposition with a unique 0-handle. Furthermore, Weinstein 1-handles can be attached in a unique way, hence the more interesting part of the handlebody decomposition is the attachment of Weinstein 2-handles along Legendrian knots in $\#^k(S^1 \times S^2)$ equipped with the unique Stein-fillable contact structures, where k is the number of 1-handles. If $k = 0$, the 2-handles are attached along Legendrian knots in S^3 equipped with its standard contact contact structure.

Following Gompf [14], a surgery diagram for X is usually presented as a front diagram in a *standard form*. In order to give a combinatorial description of the Chekanov-Eliashberg's Legendrian DG-algebra ([7, 12]) associated with a Legendrian link $L \subset \#^k(S^1 \times S^2)$ given in Gompf's standard form, Ekholm and Ng [11] described a procedure called *resolution* (analogous to [18] for resolutions of Legendrians in S^3). We give an example in Figure 1 and refer the readers to the original references [14, 11] for precise definitions.

We next recall the description of the Legendrian DG-algebra, here denoted $CE^*(L)$, that Ekholm and Ng provides coming from a resolution diagram. All our complexes are cohomological, thus we reverse the gradings from [11].

Fix a field $\mathbb{K}$. Let $L = L_1 \cup L_2 \cup \ldots L_s \subset \#^k(S^1 \times S^2)$ be a link given in the resolved form. Let $a_1, \ldots, a_n$ denote the crossings of the resolution diagram. One refers to the set $\{a_1, \ldots, a_n\}$ as the set of *external generators*. Note each a_i corresponds to a Reeb chord connecting L_v to L_w for some v and w, where L_v is the label of the under-strand and L_w



FIGURE 1. A Legendrian knot in Gompf normal form (top), Ekholm-Ng's resolution (middle), A simplifying Legendrian isotopy via Reidemeister III (bottom)

is the label of the over-strand at the crossing a_i. Thus, we can form the $\mathbf{k}$-bimodule

$$\mathcal{E} := \bigoplus_{i=1}^{n} \mathbb{K} a_i$$

where $e_w a_i e_v = a_i$ precisely if a_i corresponds to a Reeb chord from L_v to L_w.

In the interior region of each 1-handle, we also have generators. These will be referred to as the *internal generators*. Following Ekholm-Ng, we describe these generators. Let k be the number of 1-handles. For each $1 \leq l \leq k$, let o_l denote the number of strands of L entering to l-th handle. We label the strands on the left of the diagram from top to bottom by $1, \ldots, o_l$ and on the right of the diagram from bottom to top by $1, \ldots, o_l$ (note the switch in the ordering). For each 1-handle, there are infinitely many internal generators. For $1 \leq l \leq k$, these are labelled as follows:

$$c^0_{ij;l} \text{ for } 1 \leq i < j \leq o_l,$$
$$c^p_{ij;l} \text{ for } p > 0 \text{ and } 1 \leq i, j \leq o_l.$$

The generators $c^p_{ij;l}$ for $p \geq 0$ connect the i^{th} strand to j^{th} strand going to the handle labelled l, thus corresponds to a Reeb chord from some L_v to another L_w. We write

$$\mathcal{I} = \mathcal{I}_1 \cup \mathcal{I}_2 \cup \ldots \mathcal{I}_k$$

for the corresponding $\mathbf{k}$-bimodule generated by the internal generators.

Definition 7. We let $CE^*(L)$ to be the $\mathbf{k}$-algebra generated by

$$\begin{aligned}
&t_1, t_1^{-1}, \dots, t_s, t_s^{-1}\\
&a_1, a_2, \dots, a_n\\
&c^0_{ij;l},\ 1 \leq i < j \leq o_l,\ l = 1, \dots, k\\
&c^p_{ij;l},\ p > 0,\ 1 \leq i, j \leq o_l,\ l = 1, \dots, k.
\end{aligned}$$

with relations

$$\begin{aligned}
&t_v t_v^{-1} = t_v^{-1} t_v = e_v, \text{ for all } v = 1, \dots, s.\\
&t_v^{\pm} t_w^{\pm} = 0, \text{ for } v \neq w.
\end{aligned}$$

There is a differential on $CE^*(L)$ given on the external generators by

$$\partial(a_i) = \sum_{r \geq 0} \sum_{b_1, \dots, b_r} \sum_{\Delta \in \Delta(a_i; b_1, \dots, b_r)} \pm \{t_1^{-n_1(\Delta)}, \dots, t_s^{-n_s(\Delta)}, b_r, \dots, b_1\}$$

where b_q are either external generators or of the form $c^0_{ij;l}$ and n_v record the number of times the boundary of the holomorphic disk Δ passed through $*_v$, and the notation

$$\{t_1^{n_1(\Delta)}, \dots, t_s^{n_s(\Delta)}, b_r, \dots, b_1\}$$

stands for the time-ordered product where reading from right-to-left corresponds to the order of terms appearing in the boundary of Δ when it is traversed counter-clockwise. In particular, note that t_v and b_q maybe interlaced, as we do not necessarily insist that these commute. See [11, Sec. 2.4C] for the determination of the sign. [1]

It is possible to put a $\mathbb{Z}$-grading on CE^* (by making some extra choices). We refer to [11, Sec. 2.4B] for a detailed discussion of gradings on CE^*. Though, we use the negative of the grading given in [11] so that the differential increases this grading by 1. Note that the grading $|t_v| = 2r(L_v)$ where $r(L_v)$ is the rotation number.

The differential on the internal generators (see Def. (8)) satisfies $\partial(\mathcal{I}_l) \subset \mathcal{I}_l$ for all $l = 1, \dots, k$ and only depends on the number of strands passing through the 1-handle. Thus, the internal DG-algebra corresponding to a 1-handle with n strands is denoted by $\mathscr{I}_n$.

We shall next discuss the internal algebra $\mathscr{I}_n$ of Ekholm-Ng in more detail and show that it can be simplified to only include the free generators of the form $c^0_{ij;l}$ and $c^1_{ij;l}$. Hence, in particular, it is finitely generated.

2.1. **A simplification of Ekholm-Ng's internal DG-algebra.** In Ekholm-Ng [11], the internal algebra $\mathscr{I}_n$ was constructed geometrically from the study of Reeb chords internal to a 1-handle. It has the undesirable feature that it is infinitely generated. Here, we provide smaller models for the quasi-isomorphism type of $\mathscr{I}_n$. In particular, we compute

[1] As in [13], [10], we follow the convention that multiplications are read from *right-to-left*. This is not the same convention used in [11]. Throughout, our formulae are adjusted accordingly by taking the opposite DG-category (see [13, pg. 8]).

the cohomology of $\mathscr{I}_n$ for all n, and show that $\mathscr{I}_n$ is quasi-isomorphic to a finitely generated subalgebra of itself.

Note that throughout this paper the base ring is taken to be the semisimple ring $\mathbf{k}$ where the number of idempotents is given by the number of 2-handles in the surgery diagram corresponding to the components of the Legendrian link.

In order to achieve full generality, we shall first define $\mathscr{I}_n$ as a DG-algebra over a different ring

$$\mathbf{k}_n = \bigoplus_{i=1}^{n} \mathbb{K} e_i$$

As usual, n denotes the number of strands entering the 1-handle, irrespective of the 2-handles that these strands belong to. On the other hand, the internal DG-subalgebra of $CE^*(L)$ for a particular L is obtained from $\mathscr{I}_n$ by a base change $\mathbf{k}_n \to \mathbf{k}$ given by the configuration of 2-handles entering in the relevant 1-handle. (We note that in [11], $\mathbf{k}_n$ is always taken to be the ground field $\mathbb{K}$, regardless of the number of components of L, however from the perspective of Legendrian surgery, $\bigoplus_{i=1}^{n} \mathbb{K} e_i$ is the correct choice if L is not connected).

We give a formal definition as follows:

Definition 8. We define $\mathscr{I}_n$ to be a DG-category with n-objects : $X_1, \ldots, X_n$, and morphisms $\hom(\bigoplus_i X_i, \bigoplus_i X_i)$ are freely generated by morphisms in $\hom(X_i, X_j)$ given by:

$$c_{ij}^0 : 1 \leq i < j \leq n \quad \text{and} \quad c_{ij}^p : 1 \leq p, 1 \leq i, j \leq n$$

The (cohomological) grading is defined by first fixing n integers $(m_1, m_2, \ldots, m_n)$ and then defining:

$$|c_{ij}^p| = 1 - 2p + m_j - m_i.$$

The differential ∂ is determined by

$$\begin{aligned}
\partial c_{ij}^0 &= \sum_{i<r<j} (-1)^{m_r + m_j} c_{rj}^0 c_{ir}^0, \quad i < j \\
\partial c_{ij}^1 &= \delta_{ij} + \sum_{r>i} (-1)^{m_r + m_j} c_{rj}^1 c_{ir}^0 + \sum_{r<j} (-1)^{m_r + m_j} c_{rj}^0 c_{ir}^1 \\
\partial c_{ij}^p &= \sum_{0 \leq q \leq p} \sum_{r} (-1)^{m_r + m_j} c_{rj}^{p-q} c_{ir}^q, \quad p > 1
\end{aligned}$$

(c_{ij}^0 is set to be 0 for $i \geq j$, $\delta_{ij} = e_i = e_j$ if $i = j$, and 0 otherwise) and the graded Leibniz rule:

$$\partial(a_2 a_1) = \partial(a_2) a_1 + (-1)^{|a_2|} a_2 \partial(a_1).$$

We define $\mathscr{A}_n$ to be the DG-subcategory of $\mathscr{I}_n$ with the same objects $X_1, \ldots, X_n$, but morphisms in $\hom(X_i, X_j)$ are generated over $\mathbb{K}$ only by

$$c_{ij}^0 : 1 \leq i < j \leq n \quad \text{and} \quad c_{ij}^1 : 1 \leq i, j \leq n.$$

As is customary, we use the same notation for a DG-category and its total algebra. For example, $\mathscr{I}_n$ and $\mathscr{A}_n$ denote DG-algebras over $\mathbf{k}_n$. On the other hand, given a ring map $\mathbf{k}_n \to \mathbf{k}$, $\mathscr{I}_n^{\mathbf{k}}$ and $\mathscr{A}_n^{\mathbf{k}}$ denote the corresponding DG-algebras over $\mathbf{k}$.

A helpful way of thinking about these categories is to draw a circle with n points ordered in counter-clockwise direction corresponding to the objects X_i as in Figure 2. The set of generators c_{ij}^p is in bijection with homotopy classes of paths following a clockwise flow that connects i to j. The exponent p counts the number of times such paths pass from an additional marked point $*$ between n and 1. Multiplication in the category can be thought of as broken paths, and the differential corresponds to breaking the path into two paths of shorter length, with the exception that in the case of ∂c_{ii}^1, there is an additional idempotent term e_i that appears in the differential.



FIGURE 2. Visualisation of generators c_{ij}^p.

Next, let us introduce the following A_∞-algebra, whose definition is attributed to Kontsevich in [22, Rmk. 3.11].

Definition 9. We define $\mathscr{K}_n$ to be the strictly unital A_∞-category with n-objects $Z_1, \dots, Z_n$ and morphisms given by, for $i \in \mathbb{Z}/n$,

$$\hom(Z_i, Z_i) = \mathbb{K}e_i,$$
$$\hom(Z_i, Z_{i+1}) = \mathbb{K}x_i,$$

where e_i are idempotents. We again fix n integers $(m_1, m_2, \dots, m_n)$ and define gradings

$$|x_i| = 1 + m_{i+1} - m_i \text{ for } i = 1, \dots, n-1, \text{ and } |x_n| = -1 + m_1 - m_n$$

The A_∞-operations involving idempotents are determined by strict unitality. The only other products are given by:

$$\mathfrak{m}_n(x_{i-1}, \dots, x_1, x_0, x_{n-1}, \dots, x_{i+1}, x_i) = e_i, \quad \text{for} \quad i \in \mathbb{Z}/n$$

The fact that A_∞-relations are satisfied is checked easily by the following equation that holds for all $i \in \mathbb{Z}/n$:

$$\mathfrak{m}_2(e_i, \mathfrak{m}_n(x_{i-1}, \dots, x_1, x_0, x_{n-1}, \dots, x_{i+1}, x_i))$$
$$+ (-1)^{|e_i|-1}\mathfrak{m}_2(\mathfrak{m}_n(x_{i-1}, \dots, x_1, x_0, x_{n-1}, \dots, x_{i+1}, x_i), e_i) = \mathfrak{m}_2(e_i, x_i) - \mathfrak{m}_2(x_i, e_i) = 0$$

It can be shown that for $n \geq 3$, $\mathscr{K}_n$ is not formal as an A_∞-algebra, but we have

$$\mathscr{K}_2 \simeq \mathbf{k}_2\langle x_1, x_2\rangle/\langle x_2x_1 = e_1, x_1x_2 = e_2\rangle \quad |x_1| = 1 + m_2 - m_1, |x_2| = -1 + m_1 - m_2$$

Recall that cobar and bar constructions extend to the setting of A_∞-algebras. We denote these by Ω and B, respectively. (See [10, Sec. 2] for a review that is compatible with our sign conventions.)

Lemma 10. *$\mathscr{I}_n$ is isomorphic to $\Omega\mathrm{B}\mathscr{K}_n$. In particular, there exists a (strictly unital) A_∞-functor*

$$\mathfrak{f} : \mathscr{K}_n \to \mathscr{I}_n$$

which is a quasi-isomorphism.

Proof. The fact that $\mathscr{I}_n$ is isomorphic to $\Omega\mathrm{B}\mathscr{K}_n$ essentially follows by definition of Ω and B. Indeed, the A_∞-functor $\mathfrak{f}$ is the Yoneda embedding defined by sending $\mathfrak{f}(Z_i) = X_i$ at the level of objects, and on morphisms, we set for $k \geq 1$:

$$\mathfrak{f}(x_{i+k-1}, x_{i+k-2}, \ldots, x_i) = (-1)^{|c_{ij}^p|}c_{ij}^p$$

where j and p are determined uniquely by the equation $i + k = j + np$, $1 \leq j \leq n$. Here, the indices i appearing on generators x_i are considered in $\mathbb{Z}/n$. The A_∞-functor equation takes the form (using the rules [13, Eq. 4]):

$$\sum_{0\leq q\leq p}\sum_r (-1)^{|c_{ir}^q|}\mathfrak{f}(x_{j+np-1}, \ldots, x_{r+nq})\mathfrak{f}(x_{r+nq-1}, \ldots, x_i) + \partial c_{ij}^p = \delta_{ij}\delta_{1p}$$

which coincides precisely with the definition of the differential on $\mathscr{I}_n$.

Note that in [10, Sec. 2], the signs for the cobar-bar constructions are written out explicitly. It is an easy exercise to check that the signs are compatible with the signs that appear in Definition 8 of the internal algebra.

Finally, the fact that $\mathfrak{f}$ is quasi-isomorphism follows from the well known adjunction (cf. [10, Sec. 2.2.2]):

$$\hom_{DG}(\Omega\mathscr{C}, \mathscr{A}) \simeq \hom_{coDG}(\mathscr{C}, \mathrm{B}\mathscr{A})$$

where we put $\mathscr{C} = \mathrm{B}\mathscr{K}_n$ and $\mathscr{A} = \Omega\mathrm{B}\mathscr{K}_n$. □

Corollary 11. *$H^*(\mathscr{I}_n)$ is generated over $\mathbf{k}_n$ by the elements*

$$c_{12}^0, c_{23}^0, \ldots, c_{(n-1)n}^0, c_{n1}^1.$$

We next use this computation to show that the DG-subalgebra $\mathscr{A}_n$ of $\mathscr{I}_n$ generated by only the elements

$$c_{ij}^0 : 1 \leq i < j \leq n \quad \text{and} \quad c_{ij}^1 : 1 \leq i, j \leq n.$$

is quasi-isomorphic to $\mathscr{I}_n$.

Theorem 12. *There exists a retraction $\mathfrak{r} : \mathscr{I}_n \to \mathscr{A}_n$ given by*

$$\begin{aligned}\mathfrak{r}_{|\mathscr{A}_n} &= \mathrm{Id}_{\mathscr{A}_n}\\ \mathfrak{r}(c^p_{ij}) &= C_{p-1}\sum_{k_1,\dots,k_{2p-2}} (-1)^{\dagger} c^1_{k_{2p-2}j}\dots c^1_{k_1k_2}c^1_{ik_1} \quad \textit{for } p>1,\end{aligned}$$

where $C_p = \frac{1}{p+1}\binom{2p}{p}$ is the p^{th} Catalan number and $\dagger = m_{k_1}+\ldots+m_{k_{2p-2}}$, that induces a quasi-isomorphism.

Proof. Since, by Cor. (11), the cohomology is generated by cycles in $\mathscr{A}_n$, all we need to check is that $\mathfrak{r}$ is a DG-algebra map, and as $\mathscr{I}_n$ is free as an algebra, this amounts to showing that $\mathfrak{r}$ commutes with the differential on the generators.

For $p>1$,

$$\begin{aligned}\mathfrak{r}\partial(c^p_{ij}) = &\sum_{k_1,k_2,\dots,k_{2p-1}} (-1)^{\dagger+m_j+m_{k_{2p-1}}} C_{p-1}\left(c^0_{k_{2p-1}j}c^1_{k_{2p-2}k_{2p-1}}\dots c^1_{ik_1} + c^1_{k_{2p-1}j}\dots c^1_{k_1k_2}c^0_{ik_1}\right)\\ &+\sum_{k_1,k_2,\dots,k_{2p-3}} (-1)^{\dagger+m_j-m_{k_{2p-2}}}(C_{p-2}+C_2C_{p-3}+\cdots C_{p-3}C_2+C_{p-2})c^1_{k_{2p-3}j}\dots c^1_{ik_1}\end{aligned}$$

whereas

$$\begin{aligned}\partial\mathfrak{r}(c^p_{ij}) = C_{p-1}\Bigg(&\sum_{k_1,k_2,\dots,k_{2p-1}} (-1)^{\dagger+m_j+m_{k_{2p-1}}}\left(c^0_{k_{2p-1}j}c^1_{k_{2p-2}k_{2p-1}}\dots c^1_{ik_1} + c^1_{k_{2p-1}j}\dots c^1_{k_1k_2}c^0_{ik_1}\right)\\ &+\sum_{k_1,k_2,\dots,k_{2p-3}} (-1)^{\dagger+m_j-m_{k_{2p-2}}}c^1_{k_{2p-3}j}\dots c^1_{ik_1}\Bigg)\end{aligned}$$

after cancellations. The fact that Catalan numbers satisfy the recurrence relation

$$C_{p-1} = \sum_{q=0}^{p-2} C_qC_{p-q-2}$$

finishes the proof. □

Remark 13. An explicit chain homotopy $\mathfrak{H}$ between $\mathfrak{i}\circ\mathfrak{r} : \mathscr{I}_n \to \mathscr{I}_n$ and the identity map is defined by

$$\begin{aligned}\mathfrak{H}_{|\mathscr{A}_n} &= 0\\ \mathfrak{H}(c^p_{ij}) &= \sum_{q=2}^{p} C_{p-q+1}\sum_{k_1,\dots,k_{2p-2q+1}} (-1)^{\ddagger} c^q_{k_{2p-2q+1}j}c^1_{k_{2p-2q}k_{2p-2q+1}}\dots c^1_{k_1k_2}c^1_{ik_1} \quad \text{for } p>1\\ \mathfrak{H}(a_2a_1) &= \mathfrak{H}(a_2)\mathfrak{r}(a_1) - (-1)^{|a_2|}a_2\mathfrak{H}(a_1)\end{aligned}$$

where $\ddagger = m_{k_1}+m_{k_2}+\ldots+m_{k_{2p-2q+1}}+m_j$.



FIGURE 3. $\mathscr{K}_{n+1}$ represented as the partially wrapped category of $\mathbb{D}^2$ with $(n+1)$ marked points.

Remark 14. For the readers who prefer a less computational approach, we outline another proof that uses more abstract machinery but is less explicit.

The category $\mathscr{K}_{n+1}$ appears as the partially wrapped Fukaya category of the $\mathbb{D}^2$ with $(n+1)$ marked points, see [15, Sec. 3.3]. In Figure 3, one can find the Lagrangians in this category that correspond to the objects Z_i for $i = 1, \ldots, n+1$. As noted in [15, Sec. 3.3], one can easily check that the object $Z_{n+1}[-|x_{n+1}|]$ is isomorphic to the twisted complex:

$$Z_1 \xrightarrow{x_1} Z_2[|x_1| - 1] \xrightarrow{x_2} \cdots \xrightarrow{x_{n-1}} Z_n[|x_1| + |x_2| + \ldots + |x_{n-1}| + 1 - n]$$

where the isomorphism maps are given by x_n and x_{n+1}.

Thus, if we take the full subcategory of $\mathscr{K}_{n+1}$ consisting of objects $Z_1, \ldots, Z_n$ and localise it along the above twisted complex representing Z_{n+1}, we should obtain $\mathscr{K}_n$ (as shown in [15, Prop. 3.5], localising along the object Z_{n+1} corresponds to the removal of the marked point near which it is supported).

Now, it is an easy exercise in localisation of DG-categories ([9], cf. [15, Sec 3.5]) to show that localisation of $Z_1, \ldots, Z_n$ along the above twisted complex can be identified directly with $\mathscr{A}_n$.

We next show that given a ring map $\mathbf{k}_n \to \mathbf{k}$, $\mathfrak{r}$ also gives an isomorphism between the DG-algebras $\mathscr{I}_n^{\mathbf{k}}$ and $\mathscr{A}_n^{\mathbf{k}}$ over $\mathbf{k}$.

Lemma 15. *For any ring map $\mathbf{k}_n \to \mathbf{k}$, the retraction $\mathfrak{r} : \mathscr{I}_n^{\mathbf{k}} \to \mathscr{A}_n^{\mathbf{k}}$ induces a quasi-isomorphism.*

Proof. It suffices to prove the statement for $\mathbf{k} = \mathbb{K}$, and the ring map $\mathbf{k}_n \to \mathbb{K}$ given $e_i \to 1$, as any other $\mathscr{I}_n^{\mathbf{k}}$ appears as a subcomplex of $\mathscr{I}_n^{\mathbb{K}}$ that is respected by $\mathfrak{r}$. Let Δ_0 be the $\mathbf{k}_n$-bimodule generated (over $\mathbb{K}$) by $e_i \otimes_{\mathbb{K}} e_j$ for $i \neq j$. Then, we observe that

$$\mathscr{I}_n^{\mathbb{K}} \simeq \mathscr{I}_n \oplus (\mathscr{I}_n \otimes_{\mathbf{k}_n} \Delta_0 \otimes_{\mathbf{k}_n} \mathscr{I}_n) \oplus (\mathscr{I}_n \otimes_{\mathbf{k}_n} \Delta_0 \otimes_{\mathbf{k}_n} \mathscr{I}_n \otimes_{\mathbf{k}_n} \Delta_0 \otimes_{\mathbf{k}_n} \mathscr{I}_n) \oplus \cdots$$

The result then follows from Theorem 12 and the Künneth formula. □

We note that cohomology of $\mathscr{I}_n$ and $\mathscr{I}_n^{\mathbf{k}}$ can be very different. For example, in what follows we will extensively use the fact that

$$\mathscr{I}_2^{\mathbb{K}} \simeq \mathbb{K}[z, z^{-1}]$$

which is infinite-dimensional over $\mathbb{K}$. On the other hand, as we proved in Corollary 11, cohomology of $\mathscr{I}_2$ is 2-dimensional over $\mathbf{k}_2$.

3. Plumbings of cotangent bundles of 2-spheres

Throughout this section, we assume that Γ is a finite graph with $\mathbf{g} = 0$. We first construct a Legendrian link L_Γ in $\#^k(S^1 \times S^2, \xi_{std})$ associated to Γ. An important feature of this construction is that the symplectic 4-manifold obtained by adding symplectic 2-handles along L_Γ to $\#^k(S^1 \times \mathbb{D}^3, \omega_{std})$ is homotopic, as a Weinstein manifold, to the plumbing of copies of T^*S^2 with respect to the graph Γ.

After the construction, we compute the Chekanov-Eliashberg DG-algebra $\mathscr{B}_\Gamma = CE^*(L_\Gamma)$ of L_Γ and prove that it is quasi-isomorphic to the derived multiplicative preprojective algebra associated to Γ.

3.1. **A Legendrian surgery presentation of plumbings.** Choose a spanning tree T of Γ and embed it in the upper half plane $\mathbb{R} \times \mathbb{R}_{\geq 0}$ such that all the vertices lie on $\mathbb{R} \times \{0\}$. This can be easily done recursively (in multiple ways). We give an illustrating picture in Fig. (4) and leave the elaboration of the recursive argument to the reader.

Corresponding to the spanning tree T, we will have 2-handles in the Legendrian surgery picture of X_Γ. The complement of T will correspond to 1-handles.

Before we put the graph Γ in a standard form, we choose an orientation and order on the edges of Γ. Note that, as is the case for the multiplicative preprojective algebra Λ_Γ, $\mathscr{B}_\Gamma$ is also independent of these choices. Then for each arrow $h \in \Gamma \setminus T$ from v to w, we introduce a pair of arcs in the lower half plane $\mathbb{R} \times \mathbb{R}_{<0}$, one from v to the left and the other from w to the right as in Fig. (4). We assume that the arrows in T are also oriented from right to left. We also assume that in the ordering of all the edges of Γ, the edges in T are ahead of the edges in $\Gamma \setminus T$.

Once Γ is in a standard form, the endpoints of each pair of arcs corresponding to the same arrow in $\Gamma \setminus T$ is replaced by the feet of a 1-handle and each vertex v is replaced by the standard Lagrangian projection of a Legendrian unknot U_v with $tb = -1$. For each arrow a from v to w, we introduce a meridian m_a of the right half of U_w and an arc on the left half of U_v, (so that all meridians and arcs on an unknot are ordered according to the chosen order of the edges), replace a by a band from the arc on U_v to an arc on m_a, and then connect sum U_v and m_a using this band. There is no difference between the treatment of arrows in T and the rest, except that for $a \in T$, the corresponding band is planar whereas for $a \notin T$ the band goes over the associated 1-handle and it may go over/under other such bands. Note that these connected sum operations are performed directly on the Lagrangian projection without reverting back to the front projection first and then resolving afterwards (which would give a stabilisation of the Chekanov-Eliashberg DG-algebra we obtain below). Figures 4, 5 and 6 describe how the Legendrian L_Γ is constructed in a sufficiently complicated example, from which the general pattern is clear.

Now that we have a standard Lagrangian projection of L_Γ, here are the crossings:

- a pair of crossings c_a and c_a^* between $U_{s(a)}$ and $U_{t(a)}$ for each arrow a of Γ
- a self-crossing τ_v of U_v for each vertex v of Γ



FIGURE 4. The graph Γ in a standard form with respect to a chosen orientation

- a self-crossing ζ_a of $U_{s(a)}$ near the right foot of the 1-handle h_a, for each arrow a in $\Gamma \backslash T$.
- additional crossings associated to nonplanar bands in the above construction of L_Γ which go over/under each other

3.2. **Computation of $\mathscr{B}_\Gamma = CE^*(L_\Gamma)$.** The computations in this section are based on the simplification of the Chekanov-Eliashberg DG-algebra of a Legendrian link in $\#^k(S^1 \times S^2, \xi_{std})$ presented in Sec. (2.1), where the internal DG-algebra $\mathscr{I}_n$ is proved to be quasi-isomorphic to its finitely generated DG-subalgebra $\mathscr{A}_n$ as well as the A_∞-algebra $\mathscr{K}_n$.

The resolution of L_Γ to be used is the one constructed in Sec. (3.1) with the choice of orientation and base-points indicated in Fig. (6). In this resolution, for every 1-handle h_a induced by an arrow $a \in \Gamma \setminus T$, there are exactly two strands going over h_a which belong



FIGURE 5. The configuration of bands and different types of secondary crossings.

to the same component of L_Γ. Therefore the internal generators $c^0_{12;a}, c^1_{21;a}$ can be replaced by the generators of $\mathscr{I}_2^{\mathbb{K}} \simeq \mathbb{K}[z_a, z_a^{-1}]$ with grading 0 and the rest of the internal generators associated to h_a can be dropped from the presentation.

Except for $t_v^{\pm}$ and $z_a^{\pm}$, all the generators of the Chekanov-Eliashberg algebra $\mathscr{B}_\Gamma$ come from the crossings in the resolution. We use the same notation for these generators as the corresponding crossings, and list them as follows (see Fig. (6)).

- a pair of generators c_a, c_a^* with $|c_a| = |c_a^*| = 0$ for each arrow a of Γ,
- a generator τ_v with $|\tau_v| = -1$ and [2]

$$\partial\tau_v = \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v \prod_{s(a)=v,\ a\notin T} z_a \prod_{s(a)=v,\ a\in T} (e_v + c_a^* c_a)$$

  for each vertex v of Γ, where $s(a)$ and $t(a)$ denote the source and target vertices of a, respectively,
- a generator ζ_a with $|\zeta_a| = -1$ and

$$\partial\zeta_a = e_v + c_a^* c_a - z_a$$

[2]Although our sign conventions for CE^* follow [11], as we must choose the null-cobordant spin structures on the components of L_Γ, we apply the substitution $t_v \mapsto -t_v$ to their formulas (see [11, Rem2.7])



$$a_1 : v \to w_1, \quad a_2 : w_2 \to v$$
$$a_3 : v \to w_3, \quad a_4 : w_4 \to v$$
$$\partial\tau_v = -t_v z_{a_3}(e_v + c^*_{a_1}c_{a_1}) + (e_v + c_{a_2}c^*_{a_2})(e_v + c_{a_4}c^*_{a_4})$$
$$\partial\zeta_{a_4} = (e_{w_4} + c^*_{a_4}c_{a_4}) - z_{a_4}$$

FIGURE 6. A local picture showing the choice of attachments and the corresponding computation of the differential.

for each arrow a in $\Gamma\backslash T$, where $v = s(a)$,

- *secondary generators* associated to crossings between nonplanar bands in the construction of L_Γ

We call the generators $t_v^\pm$, $z_a^\pm$ and those in the first three classes above *primary generators*. In the next lemma, we show that the secondary generators are really an artefact of the projection and could be removed.

Lemma 16. *The DG-subalgebra $\mathscr{P}_\Gamma$ generated by primary generators of $\mathscr{B}_\Gamma$ is quasi-isomorphic to $\mathscr{B}_\Gamma$.*

Proof. We will prove that $\mathscr{B}_\Gamma$ and $\mathscr{P}_\Gamma$ by showing that the secondary generators can be destabilised (after suitable elementary automorphisms) without effecting the rest of $\mathscr{B}_\Gamma$. Technically, such a stable tame isomorphism argument requires the DG-algebra to be semi-free, but this does not become an issue in the current set-up since we will only use elementary automorphisms which are identity on the non-free generators $t_v^\pm$ and $z_a^\pm$.

First of all, note that every intersection point of the arcs in the lower half of the standard form of Γ (blue arcs in Fig. (4)) induces a set of 4 generators which correspond to the crossings between the associated bands in the resolution of L_Γ. We always assume that the band with the smaller slope goes under the other one. A consequence of this is that a polygon whose all vertices are induced by intersection of arcs (in the standard form of Γ) has at least 2 (Reeb-)positive vertices, hence do not contribute to the differential in $\mathscr{B}_\Gamma$. This is because at a vertex, the sign (of the quadrant in the interior of the polygon)

is negative if and only if the slope decreases while traversing the polygon in the clockwise direction. Moreover, no secondary generator appears in the differential of a secondary generator induced by another intersection or a primary generator.

In the rest of the proof we explain how to destabilise the secondary generators explicitly. The generators induced by the same intersection point in the standard form of Γ are destabilised simultaneously. This process varies slightly depending on whether the intersecting arcs end on the left or on the right of the diagram. Examples of different cases are indicated in Fig. (5). In the arguments below, the arrows in play are $a, a' \in \Gamma \setminus T$ from i, i' to j, j', resp., and the generators are denoted by p, q, r, s.

Case 1: Both arcs end on the left: The differentials of the generators are

$$\partial p = z_a q + r z_{a'}, \quad \partial q = s z_{a'}, \quad \partial r = -z_a s, \quad \partial s = 0$$

Once we apply the following elementary automorphisms successively

$$s \mapsto z_a^{-1} s, \quad q \mapsto q - z_a^{-1} r z_{a'}, \quad q \mapsto z_a^{-1} q$$

the differentials become

$$\partial p = q, \quad \partial q = 0, \quad \partial r = -s, \quad \partial s = 0$$

and p, q, r, s can be destabilised.

Case 2: Both arcs on the right: In this case, the differentials are

$$\partial p = (e_i + c_a c_a^*) q + r(e_{i'} + c_{a'} c_{a'}^*) \ , \quad \partial q = s(e_{i'} + c_{a'} c_{a'}^*) \ , \quad \partial r = -(e_i + c_a c_a^*) s \ , \quad \partial s = 0$$

We apply the elementary automorphisms

$$p \mapsto p + \zeta_a q + r\zeta_{a'} + \zeta_a s \zeta_{a'}, \quad q \mapsto q + s\zeta_{a'} \quad r \mapsto r - \zeta_a s$$

successively, making the differentials identical to those considered in the first case.

Case 3: The arcs end on different sides: The differentials are

$$\partial p = z_a q + r(e_{i'} + c_{a'} c_{a'}^*) \ , \quad \partial q = s(e_{i'} + c_{a'} c_{a'}^*) \ , \quad \partial r = z_a s \ , \quad \partial s = 0$$

In order to reduce to the first case, it suffices to apply the following elementary automorphisms successively

$$p \mapsto p + r\zeta_{a'}, \quad q \mapsto q + s\zeta_{a'}$$

□

Theorem 17. *Let Γ be a finite graph. The Chekanov-Eliashberg algebra $\mathscr{B}_\Gamma$ of the Legendrian link L_Γ and the derived multiplicative preprojective algebra $\mathscr{L}_\Gamma$ associated to Γ are quasi-isomorphic DG-algebras.*

Proof. By the previous lemma, it suffices to find a quasi-isomorphism between $\mathscr{P}_\Gamma$ and $\mathscr{L}_\Gamma$. In fact, there is a DG-algebra isomorphism $\varphi : \mathscr{P}_\Gamma \to \mathscr{L}_\Gamma$ defined on the primary generators as identity except for

$$\tau_v \mapsto \tau_v + t_v \left(\sum_{i=1}^{\iota_v} z_{a_1} \cdots z_{a_{i-1}} \zeta_{a_i} (e_v + c_{a_{i+1}} c_{a_{i+1}^*}) \cdots (e_v + c_{a_{\iota_v}} c_{a_{\iota_v}^*}) \right) \prod_{s(a)=v, \ a \in T} (e_v + c_a^* c_a)$$

where $\{a_1, \ldots, a_{\iota_v}\}$ is the (ordered) set of arrows with $s(a_i) = v$ and $a_i \in \Gamma \setminus T$. □

4. Plumbings of cotangent bundles of surfaces of positive genus

In this section, we generalise the construction and the computation in the previous section to plumbings of cotangent bundles of surfaces of arbitrary genus.

4.1. **A Legendrian surgery presentation.** In Fig. (7), we give an example illustrating how to draw a Legendrian surgery picture of a plumbing of cotangent bundles of surfaces of varying genus according to an arbitrary plumbing graph. Inside the boxes, we put a



Figure 7. Surgery diagram for plumbings of cotangent bundles of surfaces of varying genus. The drawn example shows a plumbing of two copies of T^*T^2 with one copy of T^*S^2.

standard surgery picture of the cotangent bundle of a genus g_v surface. This is described in Fig. (9). It is obtained by band summing the 2-handles of g_v copies of the standard

picture T^*T^2 given in Fig. (8) and then applying simplifying Reidemeister moves. There is a freedom in choosing the location of the band sum. Our choice makes it easier to see directly that the resulting DG-algebra is quasi-isomorphic to the derived version of the higher genus multiplicative preprojective algebra.



FIGURE 8. Legendrian surgery picture of T^*T^2

4.2. **Computation of the Chekanov-Eliashberg DG-algebra.** As shown in Lem. (16) the secondary generators induced by intersecting blue curves as in Fig. (5) can be destabilised without effecting the rest of the DG-algebra $\mathscr{B}_{\Gamma,\mathbf{g}}$. In the positive genus case, we have a new set of secondary generators induced by intersecting red and blue curves as in Fig. (7) which can be destabilised exactly as before. Another set of secondary generators $p_{v,k}, q_{v,k}, r_{v,k}$ and $s_{v,k}$, for $k = 1, \dots, g_v - 1$, indicated by quadruples (p_i, q_i, r_i, s_i) for $i = 1, 2, \dots, g-1$ in Fig. (9), have a configuration and destabilisation similar to those treated in the first case of the proof of Lem. (16). By these initial simplifications, $\mathscr{B}_{\Gamma,\mathbf{g}}$ is quasi-isomorphic to its DG-subalgebra $\mathscr{P}_{\Gamma,\mathbf{g}}$ generated by the primary generators below:

- generators $t_v^{\pm}$ for each vertex v,
- generators ζ_a, $z_a^{\pm}$ for each arrow a in $\Gamma \backslash T$,
- generators c_a, c_a^* for each arrow a in Γ,
- generators τ_v, $z_{v,i}^{\pm}$, $\xi_{v,j}$, $x_{v,j}$ for each vertex v, $i = 1, \dots, 2g_v$, and $j = 1, \dots, 4g_v$

For the last group of generators above, see Fig. (9) where the notation is simplified by using x_i instead of $x_{v,i}$, etc. Note that $z_{v,i}^{\pm}$ correspond to the generators of the internal DG-algebra $\mathscr{I}_2^{\mathbb{K}} \simeq \mathbb{K}[z^{\pm}]$ of the i^{th} 1-handle h_i in the diagram of the cotangent bundle of the surface of genus g_v in Fig. (9).

As in the genus 0 case, the DG-algebra $\mathscr{P}_{\Gamma,\mathbf{g}}$ of primary generators is supported in nonpositive gradings with τ_v, ζ_a and $\xi_{v,j}$ in grading -1 and the rest of the generators in grading 0. Note that although the generators $\xi_{v,4i-2}, \xi_{v,4i-1}$, for $i = 1, \dots, g_v$, are induced by the twists near the right feet of the corresponding 1-handles similar to the generators of type ζ_a, they can be destabilised and will not appear in the ultimate presentation of $\mathscr{B}_{\Gamma,\mathbf{g}}$.

Before we show how to destabilise the generators $\xi_{v,j}$ and $x_{v,j}$, let us describe the differential on $\mathscr{P}_{\Gamma,\mathbf{g}}$ as given by Fig. (9) where $z_i = z_{v,i}, \xi_j = \xi_{v,j}, x_j = x_{v,j}$, $g = g_v$:

$$\begin{aligned}
\partial\tau_v &= \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v x_1 \prod_{s(a)=v,\ a\notin T} z_a \prod_{s(a)=v,\ a\in T} (e_v + c_a^* c_a) \\
\partial\zeta_a &= e_{s(a)} + c_a^* c_a - z_a \\
\partial\xi_{4k-3} &= -z_{2k} x_{4k-3} + x_{4k-2} x_{4k-1} x_{4k}, \text{ for } k = 1, \dots, g-1 \\
\partial\xi_{4k-2} &= x_{4k-2} - z_{2k-1}, \text{ for } k = 1, \dots, g \\
\partial\xi_{4k-1} &= x_{4k-1} - z_{2k}, \text{ for } k = 1, \dots, g \\
\partial\xi_{4k} &= e_v - x_{4k+1} z_{2k-1} x_{4k}, \text{ for } k = 1, \dots, g-2 \\
\partial\xi_{4g-3} &= -z_{2g} x_{4g-3} + x_{4g-2} x_{4g-1} \\
\partial\xi_{4g-4} &= e_v - x_{4g-3} x_{4g} z_{2g-3} x_{4g-4} \\
\partial\xi_{4g} &= e_v - z_{2g-1} x_{4g}
\end{aligned}$$

The following lemma suffices to generalise the quasi-isomorphism between $\mathscr{B}_{\Gamma,\mathbf{g}}$ and $\mathscr{L}_{\Gamma,\mathbf{g}}$ proved in Thm. (17) for $\mathbf{g} = 0$.

Lemma 18. *The Chekanov-Eliashberg DG-algebra $\mathscr{B}_{\Gamma,\mathbf{g}}$ of the Legendrian link $L_{\Gamma,\mathbf{g}}$ associated to a graph Γ and an s-tuple $\mathbf{g}$ is quasi-isomorphic to the DG-algebra generated by*

- *generators τ_v, $t_v^{\pm}$ for each vertex v,*
- *generators ζ_a, $z_a^{\pm}$ for each arrow a in $\Gamma\backslash T$,*
- *generators c_a, c_a^* for each arrow a in Γ,*
- *generators $\alpha_{v,i}^{\pm}$, $\beta_{v,i}^{\pm}$ for each vertex v, $i = 1, \dots, g_v$*

with $|\zeta_a| = |\tau_v| = -1$, $|t_v^{\pm}| = |z_a^{\pm}| = |c_a| = |c_a^*| = |\alpha_{v,i}^{\pm}| = |\beta_{v,i}^{\pm}| = 0$ and nontrivial differentials

$$\begin{aligned}
\partial\zeta_a &= e_v + c_a^* c_a - z_a \\
\partial\tau_v &= \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \prod_{s(a)=v,\ a\notin T} z_a \prod_{s(a)=v,\ a\in T} (e_v + c_a^* c_a)
\end{aligned}$$

where $[x, y] = xyx^{-1}y^{-1}$.

Proof. We start with the presentation of $\mathscr{P}_{\Gamma,\mathbf{g}} \simeq \mathscr{B}_{\Gamma,\mathbf{g}}$ preceding the statement and (as in the proof of Lem. (16)) utilise an extended notion of stable tame isomorphism where all the elementary automorphisms are identity on the non-free generators $t_v^{\pm}, z_a^{\pm}, z_i^{\pm}$. With the help of such automorphisms, we destabilise the pairs $(\xi_{v,j}, x_{v,j})$ for all j. We do this in groups of four, starting with larger j and moving our way down to $j = 1$. Note that the treatment of the initial group is slightly different than the rest because of the unique feature of the lowest piece of Fig. (9). (In the rest of the proof, we fix v and use the simplified notation g, x_j, ξ_j and z_j for g_v, $x_{v,j}, \xi_{v,j}$, and $z_{v,j}$, respectively.)



FIGURE 9. The green part of the 2-handle of $T^*\Sigma_g$ is analogous to Fig. (6)

The first four automorphisms we apply are

$$x_{4g} \mapsto z_{2g-1}^{-1}(x_{4g} + e_v), \quad x_{4g-1} \mapsto x_{4g-1} + z_{2g}, \quad x_{4g-2} \mapsto x_{4g-2} + z_{2g-1}$$

followed by

$$\xi_{4g-3} \mapsto \xi_{4g-3} + \xi_{4g-2}(x_{4g-1} + z_{2g}) + z_{2g-1}\xi_{4g-1}, \quad \xi_{4g-4} \mapsto \xi_{4g-4} + x_{4g-3}z_{2g-1}^{-1}\xi_{4g}z_{2g-3}x_{4g-4}$$

then

$$x_{4g-3} \mapsto z_{2g}^{-1}(x_{4g-3} + z_{2g-1}z_{2g})$$

and finally

$$\xi_{4g-4} \mapsto \xi_{4g-4} + z_{2g}^{-1}\xi_{4g-3}z_{2g-1}^{-1}z_{2g-3}x_{4g-4}$$

After these automorphisms, the relevant differentials are changed as follows

$$\partial\xi_{4g} = -\, x_{4g}, \quad \partial\xi_{4g-1} = x_{4g-1}, \quad \partial\xi_{4g-2} = x_{4g-2}, \quad \partial\xi_{4g-3} = -x_{4g-3}$$
$$\partial\xi_{4g-4} = e_v - z_{2g}^{-1}z_{2g-1}z_{2g}z_{2g-1}^{-1}z_{2g-3}x_{4g-4} = e_v - [z_{2g}^{-1}, z_{2g-1}]z_{2g-3}x_{4g-4}$$

and hence we can destabilise the pairs (ξ_i, x_i) for $i = 4g-3, \ldots, 4g$.

The next four automorphisms will help destabilise the next four pairs (ξ_i, x_i) of generators and establish a pattern. These are

$$x_{4g-4} \mapsto z_{2g-3}^{-1}[z_{2g-1}, z_{2g}^{-1}](x_{4g-4} + e_v), \quad x_{4g-5} \mapsto x_{4g-5} + z_{2g-2}, \quad x_{4g-6} \mapsto x_{4g-6} + z_{2g-3}$$

followed by

$$\begin{aligned}\xi_{4g-7} \mapsto \xi_{4g-7} &+ \left(\xi_{4g-6}(x_{4g-5} + z_{2g-2}) + z_{2g-3}\xi_{4g-5}\right) z_{2g-3}^{-1}[z_{2g-1}, z_{2g}^{-1}](x_{4g-4} + e_v) \\ &- z_{2g-3}z_{2g-2}z_{2g-3}^{-1}[z_{2g-1}, z_{2g}^{-1}]\xi_{4g-4}\end{aligned}$$

then

$$x_{4g-7} \mapsto z_{2g-2}^{-1}\left(x_{4g-7} + z_{2g-3}z_{2g-2}z_{2g-3}^{-1}[z_{2g-1}, z_{2g}^{-1}]\right)$$

and

$$\xi_{4g-8} \mapsto \xi_{4g-8} + z_{2g-2}^{-1}\xi_{4g-7}z_{2g-5}x_{4g-8}$$

These automorphisms result in the following differentials

$$\partial\xi_{4g-4} = -\, x_{4g-4}, \quad \partial\xi_{4g-5} = x_{4g-5}, \quad \partial\xi_{4g-6} = x_{4g-6}, \quad \partial\xi_{4g-7} = -x_{4g-7}$$
$$\partial\xi_{4g-8} = e_v - [z_{2g-2}^{-1}, z_{2g-3}][z_{2g-1}, z_{2g}^{-1}]z_{2g-5}x_{4g-8}$$

Applying $(g-2)$ additional sets of four automorphisms analogous to the last four above, we destabilise all the pairs (ξ_i, x_i). This turns the differential of τ_v into

$$\partial\tau_v = \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v P_C \prod_{s(a)=v,\ a\notin T} z_a \prod_{s(a)=v,\ a\in T} (e_v + c_a^* c_a)$$

where P_C is the product of commutators

$$[z_2^{-1}, z_1][z_6^{-1}, z_5]\cdots[z_{2g-2}^{-1}, z_{2g-3}][z_{2g-1}, z_{2g}^{-1}]\cdots[z_7, z_8^{-1}][z_3, z_4^{-1}] \text{ if } g \text{ is even, and}$$

$$[z_2^{-1}, z_1][z_6^{-1}, z_5]\cdots[z_{2g}^{-1}, z_{2g-1}][z_{2g-3}, z_{2g-2}^{-1}]\cdots[z_7, z_8^{-1}][z_3, z_4^{-1}] \text{ if } g \text{ is odd}$$

In order to match this last differential with the expression in the statement (which makes it easier to see the quasi-isomorphism $\mathscr{B}_{\Gamma,\mathbf{g}} \simeq \mathscr{L}_{\Gamma,\mathbf{g}}$), we simply relabel the generators $z_i^{\pm}$ so that

$$\alpha_{v,i} = z_{4i-2}^{-1}, \quad \alpha_{v,j} = z_{4g-4j+3}, \quad \beta_{v,i} = z_{4i-3}, \quad \beta_{v,j} = z_{4g-4j+4}^{-1}, \quad \text{for } \ 1 \leq i \leq \frac{g+1}{2} < j \leq g$$

□

Proof of Thm. (1): With the above lemma established, we proceed as in the proof of Thm. (17) and consider the following isomorphism $\varphi : \mathscr{P}_{\Gamma,\mathbf{g}} \to \mathscr{L}_{\Gamma,\mathbf{g}}$ of DG-algebras defined to be the identity on all primary generators except for

$$\tau_v \mapsto \tau_v + t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \left(\sum_{i=1}^{\iota_v} z_{a_1} \cdots z_{a_{i-1}} \zeta_{a_i} (e_v + c_{a_{i+1}} c_{a_{i+1}}^*) \cdots (e_v + c_{a_{\iota_v}} c_{a_{\iota_v}}^*) \right) \prod_{s(a)=v, a \in T} (e_v + c_a^* c_a)$$

where $\{a_1, \ldots, a_{\iota_v}\}$ is the (ordered) set of arrows with $s(a_i) = v$ and $a_i \in \Gamma \setminus T$. □

References

[1] M. Abouzaid, A geometric criterion for generating the Fukaya category. Publ. Math. Inst. Hautes Études Sci. 112 (2010), 191–240.

[2] M. Abouzaid, I. Smith, Exact Lagrangians in plumbings. Geom. Funct. Anal. 22 (2012), no. 4, 785–831.

[3] R. Bezrukavnikov, M. Kapranov, Microlocal sheaves and quiver varieties. Ann. Fac. Sci. Toulouse Math. (6) 25 (2016), no. 2-3, 473–516.

[4] P. Boalch, Global Weyl groups and a new theory of multiplicative quiver varieties. Geom. Topol. 19 (2015), no. 6, 3467–3536.

[5] F. Bourgeois, T. Ekholm, Y. Eliashberg, Effect of Legendrian surgery. With an appendix by S. Ganatra and M. Maydanskiy, Geom. Topol. 16 (2012), no. 1, 301–389.

[6] R. Casals, E. Murphy, Legendrian Fronts for Affine Varieties. Preprint, arXiv:1610.0697,

[7] Y. Chekanov, Differential algebra of Legendrian links. Invent. Math. 150 (2002), no. 3, 441–483.

[8] W. Crawley-Boevey, P. Shaw, Multiplicative preprojective algebras, middle convolution and the Deligne-Simpson problem. Adv. Math. 201 (2006), no. 1, 180–208.

[9] V. Drinfeld, DG quotients of DG categories. J. Algebra 272 (2004), no. 2, 643–691.

[10] T. Ekholm, Y. Lekili, Duality between Lagrangian and Legendrian invariants. Preprint, arXiv:1701.01284.

[11] T. Ekholm, L. Ng, Legendrian contact homology in the boundary of a subcritical Weinstein 4-manifold. J. Differential Geom. 101 (2015), no. 1, 67–157.

[12] Y. Eliashberg, Invariants in contact topology. Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998). Doc. Math. 1998, Extra Vol. II, 327–338.

[13] T. Etgü, Y. Lekili, Koszul duality patterns in Floer theory. Geom. Topol. (to appear), arXiv:1502.07922.

[14] R. Gompf, Handlebody construction of Stein surfaces. Ann. of Math. (2) 148 (1998), no. 2, 619–693.

[15] F. Haiden, L. Katzarkov, M. Kontsevich, Flat surfaces and stability structures. Preprint, arXiv:1409.8611.

[16] A. Keating, Lagrangian tori in four-dimensional Milnor fibres. Geom. Funct. Anal. 25 (2015), no. 6, 1822–1901.

[17] D. Nadler, E. Zaslow , Constructible sheaves and the Fukaya category. J. Amer. Math. Soc. 22 (2009), no. 1, 233–286.

[18] L. Ng, Computable Legendrian invariants. Topology 42 (2003), no. 1, 55–82.

[19] T. Pantev, B. Toën, M. Vaquié, G. Vezzosi, Shifted symplectic structures. Publ. Math. Inst. Hautes Études Sci. 117 (2013), 271–328.
[20] J. Pascaleff, Floer cohomology in the mirror of the projective plane and a binodal cubic curve. Duke Math. J. 163 (2014), no. 13, 2427–2516.
[21] P. Seidel, Graded Lagrangian submanifolds. Bull. Soc. Math. France 128 (2000), no. 1, 103–149.
[22] P. Seidel, Fukaya categories and Picard-Lefschetz theory. Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008.
[23] V. Shende, A. Takeda, Symplectic structures from topological Fukaya categories, Preprint, arXiv:1605.02721.
[24] D. Yamakawa, Geometry of multiplicative preprojective algebra. Int. Math. Res. Pap. IMRP 2008, Art. ID rpn008, 77pp.
[25] W-K Yeung, Relative Calabi-Yau completions. Preprint, arXiv:1612.06352.

Department of Mathematics, Koç University, Sariyer, Istanbul 34450 TURKEY
Department of Mathematics, Princeton University, Princeton, NJ 08540 USA
E-mail address: tetgu@ku.edu.tr, tetgu@math.princeton.edu

King's College London, Strand, London, UK
E-mail address: yanki.lekili@kcl.ac.uk